



GENERAL HEALTH TESTING

Ferritin Information Sheet

Iron and Ferritin

Iron is a mineral with several essential functions in the body. It is required to transport oxygen around the body and is also essential for growth, normal cell function, and the production of connective tissue and some hormones (1). Approximately 25% of the iron in a normal adult is present in a storage form (2), with the most common form being ferritin (accounting for about 2/3 of storage iron) (3).

Ferritin analyses

Ferritin analyses provide a sensitive, specific, and reliable measurement for determining iron deficiency at an early stage (4). They are useful for monitoring the reaccumulation of iron stores in iron-deficient individuals who are taking iron supplements. Ferritin analyses are also beneficial for determining iron overload and response to iron chelating agents (5).

What are optimal ferritin levels?

Healthy ferritin levels are 40 – 300 ng/mL for males and 20 – 200 ng/mL for females (6). Ferritin levels below 10 ng/mL are indicative of iron deficiency anemia, while levels above 200-300 ng/mL may indicate hemochromatosis or other health complications.

Ferritin Information Sheet

What are the signs of iron deficiency?

Low iron levels inhibit the production of hemoglobin, resulting in reduced red blood cells and a condition called anemia, which affects an estimated two billion people around the globe (7). Symptoms include:

- Tiredness
- Fatigue
- Pale skin
- Shortness of breath
- Headaches
- Dizziness

These initial symptoms of deficiency can go unnoticed, but if left untreated, anemia can have serious repercussions, including impaired cognitive function, disturbances in the digestive system, and impaired immunity. Pregnant women, young children and frequent blood donors are at a much higher risk of iron deficiency (8).

What are the signs of excess iron?

Increased iron concentrations occur in hemochromatosis and acute liver disease (9). Excess iron cannot be naturally excreted from the body, so it accumulates in organs and tissues, eventually causing serious health complications.

The symptoms of iron overload include:

- Fatigue
- Joint pain
- Abdominal pain
- Memory problems
- Depression
- Decreased sex drive
- Shortness of breath
- Heart flutters

Further serious complications can occur in untreated individuals, including heart failure, liver cirrhosis and disease, and endocrine problems (10).

Where can I find more info?

Visit www.genetrackdiagnostics.com for full test information, including specimen collection requirements

CONTACT US:

Email: support@genetrackdiagnostics.com

Phone: 1-888-802-0703

NOTE:

This brochure is provided for general information purposes only. It is not intended to replace medical advice from a health professional.

References:

- (1) Bothwell TH, et al. (1989). Nutritional iron requirements and food iron absorption. *J Int Med.* 226(5), 357-365.
- (2) Krause JR, Stolc V. (1979). Serum Ferritin and Bone Marrow Iron Stores, Correlation with Absence of Iron in Biopsy Specimens. *Am J Clin Pathol.* 72, 817-820.
- (3) Skikne BS, Cook JD. (1981). Serum Ferritin in the Evaluation of Iron Status. *Lab Management.* 19, 31-35.
- (4) Bates HM. (1980). How to Detect Iron Deficiency Before Anemia Develops. *Laboratory Pathfinder.* Jan 1980:17-22.
- (5) Lipschitz DA, Cook JD, Finch CA. (1974). A Clinical Evaluation of Serum Ferritin as an Index of Iron Stores. *N Engl J Med.* 290, 1213-1216.
- (6) Camaschella C. Iron-deficiency anemia. *N Engl J Med.* 2015 May 7;372(19):1832-43.
- (7) Zimmermann MB and Hurrell RF. (2007). Nutritional iron deficiency. *The Lancet.* 370(9586), 511-520.
- (8) Camaschella C. (2015). Iron-Deficiency Anemia. *N Engl J Med.* 372, 1832-1843.
- (9) Witte DL, et al. (1996). Hereditary hemochromatosis. *Clinica Chimica Acta.* 245(2), 139-200.
- (10) Beutler E, Felitti V, Gelbart T, Ho N. (2001) Genetics of Iron Storage and Hemochromatosis. *Drug Metab Dispos.* 29(4):495-499.