



HORMONE TESTING

Free Testosterone Information Sheet

What is Testosterone?

Testosterone is the primary sex hormone in males, but it is also important in females. Testosterone is secreted from the testicles in males, and from the ovaries in females.

What are the roles of testosterone?

Testosterone has an important role during the life cycle of both males and females. Testosterone levels before birth are associated with sex formation (1), potentially brain masculinization during early infancy, and pubertal effects, including penis or clitoris enlargement, increased libido, remodelling of facial bones, acne, pubic hair, and deepening of voice (2).

In adult males, testosterone is necessary for normal sperm development, enhances muscle growth, and regulates platelet aggregation (3). In females, testosterone also enhances muscle and bone growth, may modulate the physiology of vaginal tissue, and contributes to female genital sexual arousal (4).

Why measure free testosterone?

In the circulation, approximately 97% of testosterone is transported by proteins, in particular the sex hormone-binding globulin (SHBG), and also weakly bound to albumin (5). Only the remaining small fraction of testosterone is circulating in the free form, and it is this free testosterone that is considered the active fraction.

What is the link between testosterone and metabolism?

Variations in testosterone levels also influence metabolic parameters, particularly in the control of glucose and energy metabolism (6). In males, low serum testosterone levels are associated with an adverse metabolic profile, including an increased risk of insulin resistance (7), obesity and type 2 diabetes (8).

In contrast, high free testosterone levels in females are associated with increased risk of elevated BMI, insulin resistance and type 2 diabetes (9)

Free Testosterone Patient Brochure

What are the signs of low testosterone in males?

Symptoms of low testosterone can include:

- Low sex drive
- Fatigue
- Reduced muscle mass
- Irritability
- Erectile dysfunction
- Infertility
- Weight gain
- Osteoporosis
- Depression

Low testosterone can occur due to conditions that males are born with, such as Klinefelter syndrome, or testosterone may also decline for other reasons, including testicle injury, chemotherapy, infection, autoimmune disease, malnutrition, and hormonal imbalances.

Testosterone therapy is available for men with low testosterone levels, although often other changes, such as more physical exercise for overweight men, can increase testosterone levels (10).

What are the signs of high testosterone in males?

Elevated testosterone in males is not a common problem, and is most often observed in athletes who use anabolic steroids, testosterone, or related hormones to increase muscle mass and athletic performance. Elevated testosterone can cause many health complications (11), including:

- Low sperm counts
- Impotence
- Liver disease
- Acne
- Blood clots
- Aggressive behaviour
- Damage to heart tissue
- Weight gain
- Mood swings
- Excessive body hair
- Sleep apnea

Where can I find more info?

Visit www.genetrackdiagnostics.com for full test information, including specimen collection requirements

CONTACT US:

Email: support@genetrackdiagnostics.com
Phone: 1-888-802-0703

What are the signs of abnormal testosterone in females?

The signs of reduced testosterone in females include low muscle mass, poor sleep, lack of energy, depression, low libido, and vaginal dryness. Excess testosterone can lead to acne, polycystic ovary syndrome, irregular menstruation, increased body hair, and low HDL (good) cholesterol.

References:

- (1) Swaab DF, & Garcia-Falgueras A. (2009). Sexual differentiation of the human brain in relation to gender identity and sexual orientation. *Funct Neurol*, 24 (1), 17-28.
- (2) Raggatt LJ, & Partridge NC. (2010). Cellular and molecular mechanisms of bone remodeling. *J Biol Chem*, 285 (33), 25103-25108.
- (3) Ajayi AA, Mathur R, & Halushka PV. (1995). Testosterone increases human platelet thromboxane A2 receptor density and aggregation responses. *Circulation*, 91 (11), 2742-2747.
- (4) Traish AM, et al. (2002). Role of androgens in female genital sexual arousal: receptor expression, structure, and function. *Fertil Steril*, 77 (Suppl 4), S11-S18.
- (5) Dunn JF, Nisula BC, & Rodbard D. (1981). Transport of Steroid Hormones: Binding of 21 Endogenous Steroids to Both Testosterone-Binding Globulin and Corticosteroid-Binding Globulin in Human Plasma. *J Clin Endocrin Metab*, 53, 58-68.
- (6) Navarro G, et al. (2015). The role of androgens in metabolism, obesity, and diabetes in males and females. *Obesity*. 23(4).
- (7) Pitteloud N, et al. (2005). Relationship Between Testosterone Levels, Insulin Sensitivity, and Mitochondrial Function in Men. *Diabetes Care*. 28(7), 1636-1642.
- (8) Dhindsa S, et al. (2010). Testosterone Concentrations in Diabetic and Nondiabetic Obese Men. *Diabetes Care*. 33(6), 1186-1192.
- (9) Kim SY, et al. (1993). Direct Relationship between Elevated Free Testosterone and Insulin Resistance in Hyperprolactinemic Women. *Korean J Intern Med*. 8(1), 8-14.
- (10) What is Low Testosterone? Urology Care Foundation.
- (11) Testosterone – What It Does And Doesn't Do. (2019). Harvard Health Publishing, Harvard Medical School.

NOTE:

This brochure is provided for general information purposes only. It is not intended to replace medical advice from a health professional.